

WO 01/05752

1

PCT/FR00/01579

AMINOGUANIDINE BICARBONATE WITH PARTICULAR PROPERTIES
AND PROCESS FOR MANUFACTURING IT

The present invention relates to a process
5 for manufacturing aminoguanidine bicarbonate. The
invention also relates to an aminoguanidine bicarbonate
with particular properties.

The manufacture of aminoguanidine bicarbonate
(AGB) by reacting an aqueous solution of cyanamide with
10 hydrazine followed by an addition of CO₂ is known. Since
the placing in contact of cyanamide with hydrazine in
alkaline medium also leads to the dimerization of the
cyanamide, it is necessary to use a large excess of
cyanamide to achieve a suitable yield of aminoguanidine
15 bicarbonate.

Thus, patent DD 689 191 teaches working with
a 100% excess of cyanamide (i.e. cyanamide/hydrazine
molar ratio = 2/1) to obtain a yield (relative to the
hydrazine used) of 80% of AGB, after reaction for 60
20 hours. An AGB yield of about 90% may be achieved after
reaction for 27 hours when concentrated solutions of
cyanamide and hydrazine are used with a 100% excess of
cyanamide (DD 730 331).

Since cyanamide is a very expensive product,
25 attempts to reduce this excess have been the subject of
much research.

206240" ETT E001

Specifically, patent SU 981 314 discloses a cyanamide/hydrazine molar ratio of between 1.25 and 1.8. It is mentioned that an AGB yield (relative to the hydrazine) of 95% is obtained with a

5 cyanamide/hydrazine molar ratio of 1.8. It also teaches that the yields fall to 90% and 85% for molar ratios of 1.5 and 1.25, respectively.

The same trend has been observed by other authors. Thus, a fall of about 12 points in the yield
10 was recorded when the cyanamide/hydrazine molar ratio went from 1.2 to 1 (DD 249 009).

The combined literature in this field encourages a person skilled in the art to work with an excess of cyanamide to obtain a yield of aminoguanidine
15 bicarbonate.

The Applicant Company has developed a process for manufacturing aminoguanidine bicarbonate from cyanamide and hydrazine and has observed, surprisingly, by working with a slight deficit of cyanamide relative
20 to the stoichiometric amount, yields of AGB that are as high as, or even higher than, those achieved with processes using a large excess of cyanamide.

According to the present invention, the process consists in reacting an aqueous solution of
25 cyanamide with an aqueous solution of hydrazine hydrate in the presence of CO_2 , characterized in that the

206240" ETT E001

process is performed with a slight deficit of cyanamide relative to the stoichiometry.

The cyanamide/hydrazine molar ratio used is preferably between 0.80 and 0.99 and advantageously
5 between 0.85 and 0.95.

The pH of the reaction medium is generally between 6.5 and 8, preferably between 7 and 7.3. The pH may be adjusted by any suitable means and especially using CO₂.

10 The concentration of the aqueous solutions may vary within a wide range. It is usually preferred to use an aqueous cyanamide solution with a concentration of between 15% and 50% by weight. The hydrazine concentration in the aqueous solution is
15 advantageously between 15% and 64% by weight.

The temperature of the reaction medium is generally between 35°C and 70°C. A temperature of between 40°C and 50°C gives an aminoguanidine bicarbonate whose structure and specific properties are
20 commercially very advantageous.

One embodiment consists in adjusting, using CO₂ (carbon dioxide), the pH of the hydrazine hydrate solution to the desired value and then in introducing an aqueous solution of cyanamide once the temperature
25 of the hydrazine solution has been raised to about a few degrees below the temperature chosen for the reaction.

The pH of the reaction medium is maintained using CO₂ at the desired value during the introduction, or addition, of the cyanamide solution and throughout the reaction.

5 Another embodiment consists in simultaneously adding an aqueous hydrazine hydrate solution and carbon dioxide to an aqueous cyanamide solution initially maintained a few degrees below the temperature chosen for the reaction.

10 Irrespective of the embodiment, the total reaction time is generally between 6 and 15 hours and preferably between 7 and 10 hours. The duration of the addition of cyanamide or of hydrazine hydrate is generally between 1 and 3 hours and preferably in the
15 region of 2 hours.

After the reaction, the reaction medium is cooled to room temperature and the aminoguanidine bicarbonate thus obtained is spin-filtered or filtered and optionally dried.

20 With the process according to the invention, yields of greater than 90% and preferably greater than 95% are obtained with a purity of greater than 99%, or even greater than 99.5%.

25 A subject of the present invention is also an aminoguanidine bicarbonate with a particular structure and particular specific properties. It is characterized by a virtually spherical crystal aggregate with a mean

2003113-042902

diameter of between 80 and 500 μm . The aggregate preferably has a mean diameter of between 100 and 250 μm , the mean diameter being determined by laser granulometry.

5 The aminoguanidine bicarbonate according to the invention also has the advantage of being easy to separate out from the reaction medium by any known means, for example by filtering or spin-filtering and drying, thus being distinguished from platelet
10 crystals.

EXPERIMENTAL SECTION

Example 1

110.9 g of hydrazine hydrate with a purity of 99.2% (2.2 mol) and 300 g of demineralized water are
15 placed in a one liter reactor at room temperature. The pH of the aqueous solution is in the region of 11. Carbon dioxide is then bubbled into the aqueous solution for about 1 hour, which represents 58 g or 1.3 mol of CO_2 , until a pH in the region of 7 is
20 obtained, while maintaining the temperature of the solution at about 40°C.

171.4 g of an aqueous 49% cyanamide solution (2 mol) are then added over about 2 hours, while continuing the addition of CO_2 so as to maintain the pH
25 of the reaction medium in the region of 7. During the addition, the temperature of the medium is raised to 45°C and the medium is maintained at this temperature

for 8 hours with adjustment of the pH to a value in the region of 7 by small additions of CO₂.

The total amount of CO₂ added is 104 g, i.e. 2.36 mol.

5 At the end of the reaction, the reaction medium is allowed to cool to room temperature and the AGB crystals are then filtered off and washed with 250 ml of water and finally dried under vacuum at a temperature of between 35°C and 40°C.

10 After drying, 260 g of crystals with a purity of 99.7%, determined by assaying with perchloric acid, are obtained. The crude yield relative to the cyanamide is 95.6%.

The crystals obtained are in the form of
15 virtually spherical aggregates (photograph No. 1 by scanning electron microscopy).

Example 2

The process is performed as described in Example 1, except that the aqueous hydrazine hydrate
20 solution is maintained at 55°C instead of 40°C and that, during the addition of the cyanamide, the reaction medium is brought to 65°C and is maintained at this temperature for 4 hours.

After drying, 261.1 g of crystals in the form
25 of platelets (photograph No. 2) are obtained ^{with} in a purity of 99.6%. The crude yield relative to the cyanamide is 96%.

Example 3

Example 1 is extrapolated to the industrial scale, using a 15 m³ reactor.

After spin-filtering for 20 minutes, the aggregates contain a moisture content of only 7%. At the end of the spin-filtering, the aggregates are virtually spherical of the type in Example 1, with a narrow particle size distribution free of fine particles, of less than 40 µm in diameter.

Example 4

Example 2 is repeated on the industrial scale, using a 15 m³ reactor.

After spin-filtering for 3 hours, the platelets contain a moisture content of 20% and, at the end of the spin-filtering, the platelets have a mean diameter of 70 µm with a very broad particle size distribution with 20% of the population of particles having a diameter of less than 20 µm.

Example 5

The procedure described in Example 1 is repeated, except that the duration of addition of the cyanamide is 5 hours instead of 2 hours and the duration of the reaction after the addition is reduced from 8 to 5 hours.

The yield and also the purity of the AGB crystals obtained are similar to those of Example 1. However, the crystals are rather in the form of

platelets (photograph No. 3) and the spin-filtering time is longer.

Example 6

171.4 g of an aqueous 49% cyanamide solution
5 (2 mol) and 300 g of water are placed in a one liter reactor at room temperature. The pH of the resulting solution is in the region of 5. The solution is then brought to 40°C, after which 110.9 g of 99.2% hydrazine hydrate (2.2 mol) and 75 g (1.7 mol) of CO₂, to maintain
10 the pH at about 7, are simultaneously added over 2 hours. The reaction medium is then maintained at 45°C for 8 hours with a small addition of CO₂ to adjust the pH to about 7. The total amount of CO₂ added is 94.5 g (2.15 mol).

15 The reaction medium is then allowed to cool to room temperature and the AGB is filtered off and washed with 250 ml of water. Finally, it is dried under vacuum at a temperature of between 35°C and 40°C.

After drying, 259 g of AGB aggregates similar
20 to those of Example 1, with a purity of 99.6%, are obtained.

The crude yield of aminoguanidine bicarbonate is 95.2% relative to the cyanamide.